

The prevalence and molecular characterisation of blood parasites infecting the vulnerable Tamarugo Conebill (*Conirostrum tamarugense*) and other birds in the Pampa del Tamarugal, Chile

Javier Martínez^{A,D}, Rodrigo A. Vásquez^B, Alberto Marqués^A, Alazne Díez-Fernández^C and Santiago Merino^B

^ADepartamento de Biomedicina y Biotecnología, Área Parasitología, Facultad de Farmacia, Universidad de Alcalá, Alcalá de Henares, E-28871 Madrid, Spain.

^BInstituto de Ecología y Biodiversidad, Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile, Las Palmeras 3425, Ñuñoa, Santiago, Chile.

^CDepartamento de Ecología Evolutiva, Museo Nacional de Ciencias Naturales – Centro Superior de Investigaciones Científicas, J. Gutiérrez Abascal 2, E-28006 Madrid, Spain.

^DCorresponding author. Email: francisco.martinez@uah.es

Abstract. Blood parasites exert significant selective pressure, which can alter population dynamics, and the introduction of new parasite lineages in a region could lead to the extinction of endemic avian species that have not coevolved with them. The avifauna of the Pampa del Tamarugal in the desert of Atacama, Chile, is highly depauperate, with two species accounting for the greatest numbers: the widespread Rufous-collared Sparrow (*Zonotrichia capensis*) and the endemic Tamarugo Conebill (*Conirostrum tamarugense*). We used molecular and microscopic methods to estimate the prevalence of haemoparasites in birds inhabiting the Pampa del Tamarugal. The molecular screening of the samples from the most common species indicated that the Tamarugo Conebill was only infected by parasites of the genus *Leucocytozoon* (prevalence 3.5%) and the Rufous-collared Sparrow was uninfected. Only two other bird species (of a total of seven) were infected, the House Sparrow (*Passer domesticus*), by *Plasmodium relictum* (prevalence 30%), and the Eared Dove (*Zenaida auriculata*), by *Haemoproteus multipigmentatus* (prevalence 100%). The occurrence of House Sparrows infected with the *Plasmodium* haplotype GRW4 (involved in Hawaiian bird extinctions) may entail the risk of potential host-switching, something particularly relevant for the conservation of the endemic and vulnerable Tamarugo Conebill.

Additional keywords: avian malaria, haplotype GRW4, host-switching, House Sparrow, *Plasmodium*, *Leucocytozoon*.

Received 2 June 2015, accepted 5 February 2016, published online 13 April 2016

Introduction

The strong negative effects of parasitism have been reported in both managed and wild populations of organisms (Watson 2013). In birds, parasitism can have adverse effects on clutch-size, hatching and fledging success, and adult survival. These detrimental effects have been related to the presence of both ectoparasites and endoparasites (Merino and Potti 1995; Merino *et al.* 2000). Among avian parasites, blood parasites like haemosporidians (e.g. *Plasmodium*, *Haemoproteus* and *Leucocytozoon*) have been widely studied in birds since species belonging to this protozoan group are quite prevalent worldwide (Clark *et al.* 2014) and some of them are highly pathogenic depending on the particular host–parasite system (Valkiūnas 2005).

Haemosporidian parasites have heteroxenous life-cycles and are transmitted by dipteran vectors whose larvae are aquatic.

Thus, environmental constraints related to rainfall determine the incidence of these horizontally transmitted parasitic diseases (Sehgal 2015). Horizontal disease transmission has been related to higher virulence, especially if generalist parasites infect novel hosts (Combes 2005). In fact, parasite host-switching events have been reported with dramatic consequences for species without previous contact with these parasites. The role of avian malaria introduced with non-native bird species in the decline and extinction of native Hawaiian birds is a paradigmatic example (van Riper *et al.* 1986). The *Plasmodium* haplotype GRW4 seems to be solely responsible for these extinctions. The possibility of introduced host species functioning as disease reservoirs has been postulated as a serious threat to native birds in New Zealand (Tompkins and Gleeson 2006), on Sao Miguel Island, the Azores (Hellgren *et al.* 2011) and on Robinson Crusoe Island, Chile (Martínez *et al.* 2015).

Surveys on avian haemosporidians are biased towards European and North American passerines (Clark *et al.* 2014). Some studies have been performed in South America (Durrant *et al.* 2006; González *et al.* 2015), including Chile (Merino *et al.* 2008). However, there are no studies of haemoparasites infecting birds in Pampa del Tamarugal, a national reserve located in the Tarapacá region of Chile. This reserve, characterised by Tamarugo (*Prosopis tamarugo*) forest, is in the Atacama Desert, the most arid desert in the world. Owing to extreme climatic conditions (i.e. high thermic amplitude and very low rainfall), biodiversity is very low, and only 17 species of bird have been recorded (Estades 1995). The Rufous-collared Sparrow (*Zonotrichia capensis*), Tamarugo Conebill (*Conirostrum tamarugense*) and Scaly-throated Earthcreeper (*Upucerthia dumetaria*) are the most abundant species (Estades 1995).

The Tamarugo Conebill is considered endemic to the deserts of northern Chile and southern Peru (López-Calleja and Estades 1996). It is a vulnerable insectivorous bird (BirdLife International 2012) whose conservation seems to be tightly linked to management of Tamarugo forests, which support the Conebill's critical food resource, *Leptotes trigemmatum* caterpillars (López-Calleja and Estades 1996), especially during spring. Populations of Tamarugo Conebills winter at high altitudes in southern Peru (i.e. Arequipa, Tacna and Moquegua) where arid plants of the genera *Gynoxys* and *Polylepis* are common at 3400–4050 m above sea level (asl).

The Tamarugo Conebill can be considered a restricted-range species because its breeding range is small, estimated at only 108 km² (Estades 1996). Its small breeding range combined with the low avian diversity and extreme climatic conditions in that range may have allowed it to avoid some parasite lineages in the past. In addition, the high-altitude climatic characteristics of its winter range in southern Peru may also have limited encounters with some parasites transmitted by mosquitoes, such as *Plasmodium*, since optimal conditions for the reproduction of these vectors are lacking (LaPointe *et al.* 2010). Thus, in the present study, we have estimated the occurrence of haemoparasites (i.e. *Plasmodium*, *Haemoproteus*, *Leucocytozoon*, *Trypanosoma* and microfilariae) in the avian community of the Pampa del Tamarugal arid region to identify potential parasitic threats to the vulnerable Tamarugo Conebill.

Materials and methods

Capture and sampling of birds was conducted within or near the Pampa del Tamarugal National Reserve for 7 days during January 2012 (12–20 January) at three sites: La Huayca (3 days) and Canchones (2 days) within the National Reserve, and Pica, 20 km east of Pampa del Tamarugal (2 days) (see section S1 in supplementary material available online). During January mean maximum and minimum temperatures are slightly above 30°C and 10°C, respectively, and rainfall is very low, if any (Arenas 2013; Hajek and Di Castri 1975).

The Pampa del Tamarugal is a vast plain within the Atacama Desert, located 70–100 km east of the city of Iquique, at ~1000 m asl. Some areas are vegetated with *Prosopis tamarugo*, a flowering tree species belonging to the family Fabaceae. This tree species grows without rainfall and obtains water via a root-system that reaches deep ground-water supplies and possibly also obtains

water from dew. The three sampling sites were selected based on previously observed bird activity (see section S1).

The birds were captured in mist-nets, which were checked at 5-min intervals to avoid undue stress or overheating of birds. All mist-netting was done between 1000 and 1215 hours. Approximately 20 µL of blood was extracted by puncture of the tarsal or wing vein with a 0.5-mm gauge needle for blood smears and molecular analyses. Blood smears were done in the field at the time of blood extraction (see Merino *et al.* 1997) and blood for molecular analyses were stored on Whatman FTA cards (GE Healthcare Life Sciences, Little Chalfont, UK) to preserve biological samples for downstream DNA analysis. Genomic DNA was extracted from samples stored on FTA cards according to Martínez *et al.* (2009). Details of the screening of parasites by microscopy and PCR (Tables S1 and S2) are available in section S2 of the supplementary material.

Results

The numbers and species of birds captured and examined at the three sampling sites, and the number and proportion of infected birds are shown in Table 1. The most commonly captured bird species were the Tamarugo Conebill and Rufous-collared Sparrow, which together comprised >90% of the individuals captured. Overall, the prevalence of haemoparasites was low in the present survey (10 infected of 197 individuals examined = 5% (95% confidence level 2–8.1%)). Infections were found in three Tamarugo Conebills, with *Leucocytozoon* (3 of 89 examined = 3.4% (0–7.1%)); three House Sparrows, with *Plasmodium* (3 of 10 examined = 30% ((1.6–58.4%)); and four Eared Doves, with *Haemoproteus* (4 of 4 examined = 100%). No mixed infections or infections by other blood parasites were detected (Table 1).

The haplotypes isolated from the Tamarugo Conebills (GenBank accession KT363871, 719 bp), House Sparrows (GenBank accession KT363870, 824 bp) and Eared Doves (GenBank accession KT363869, 326 bp) correspond to *Leucocytozoon* sp., *Plasmodium relictum* and *Haemoproteus multipigmentatus*, respectively, as indicated by the BLAST analyses (Table 2). Lineage names, genetic identities, query cover, E value and other hosts from which these haplotypes were previously isolated are detailed in Table 2.

Microscopic examinations of blood smears, corresponding to PCR positive samples, were negative for *Plasmodium* and *Leucocytozoon* but positive for *Haemoproteus*.

Discussion

In this study, we present preliminary data on the prevalence of haemoparasites in the avian community inhabiting the Pampa del Tamarugal. Tamarugo Conebills and Rufous-collared Sparrows together comprised >90% of the individual birds captured, with other bird species scarce in the community. Overall, the prevalence of haemoparasites was low (5.08%). However, a more exhaustive sampling effort should be conducted to obtain adequate sample sizes for all species to enable an accurate estimation of overall prevalence. Nevertheless, we can provide a reasonable estimate of haemoparasite prevalence for the two most abundant bird species in the region: Rufous-collared Sparrows were uninfected, and only three of 89 Tamarugo Conebills were infected (prevalence 3.37%), with parasites of the genus *Leucocytozoon*.

Table 1. Numbers of each species captured and examined at the three sampling sites, and number and proportion (%) of infected birds

Under Number examined, values in parentheses are the proportion of individuals of a species as a percentage of the total of all birds captured ($n = 197$). Under Number infected, values in parentheses are the number infected as a proportion of the number of that species captured

Common name	Scientific name	Number captured and examined (proportion of total captures)	Number of infected birds (proportion of captures of each species infected)	Number captured at each sampling site		
				Huayca	Canchones	Pica
House Sparrow	<i>Passer domesticus</i>	10 (5.08%)	3 (30.0%) ^A	0	1	9
Eared Dove	<i>Zenaida auriculata</i>	4 (2.03%)	4 (100%) ^B	0	0	4
Tamarugo Conebill	<i>Conirostrum tamarugense</i>	89 (45.2%)	3 (3.37%) ^C	33	19	37
Cinereous Conebill	<i>Conirostrum cinereum</i>	5 (2.54%)	0	2	0	3
Rufus-collared Sparrow	<i>Zonotrichia capensis</i>	87 (44.1%)	0	20	56	11
Vermilion Flycatcher	<i>Pyrocephalus rubinus</i>	1 (0.51%)	0	0	1	0
Scaly-throated Earthreeper	<i>Upucerthia dumetaria</i>	1 (0.51%)	0	0	1	0
Total		197	10 (5.08%)	55	78	64

^AAll infected with *Plasmodium* and captured in Pica.

^BAll infected with *Haemoproteus* and captured in Pica.

^CAll infected with *Leucocytozoon*, two captured in Pica, the other in Canchones.

Table 2. Identification of amplicons detected in the present study using BLAST analysis

Amplicons in this study	Genus	Lineage	Identity	Query cover	E value	Hosts
<i>Conirostrum tamarugense</i> (GenBank KT363871; 719 bp)	<i>Leucocytozoon</i>	L-T087 or TROAED06 ^D	100%	66%	0.0	Cinereous Conebill (<i>Conirostrum cinereum</i>), Junin, Peru, (GenBank JQ988442); Buff-browed Foliage-gleaner (<i>Syndactyla rufosuperciliata</i>), Cuzco, Peru (GenBank JQ988495); Rufous-breasted Chat-tyrant (<i>Ochthoeca rufipectoralis</i>), Junin, Peru (GenBank JQ988532); House Wren (<i>Troglodytes aedon</i>), Peru (GenBank KF767441 and KF767437)
<i>Passer domesticus</i> (GenBank KT363870; 824 bp)	<i>Plasmodium</i>	GRW04 ^A or P15	100%	100%	0.0	Hawai'i 'amakihi (<i>Hemignathus virens</i>), Hawaii (GenBank AY733090); Great Reed-Warbler (<i>Acrocephalus arundinaceus</i>), Sweden (GenBank AF254975); House Sparrow (<i>Passer domesticus</i>), USA (GenBank AY172850); <i>Culex pipiens quinquefasciatus</i> , Japan (GenBank AB308050)
<i>Zenaida auriculata</i> (GenBank KT363869; 326 bp)	<i>Haemoproteus</i>	SocH6	100%	99%	1e-167	Mourning Dove (<i>Zenaida macroura</i>), Socorro Island, Mexico (GenBank JN788937)

^ALineage TROAED06 and GRW04 from MalAvi database.

The three Tamarugo Conebills were PCR positive for the parasite but smears were negative, indicating a very low parasitaemia. The obtained haplotype seems to be generalist since it was previously detected in other species of birds, such as the Cinereous Conebill (*Conirostrum cinereum*), Buff-browed Foliage-gleaner (*Syndactyla rufosuperciliata*), Rufous-breasted Chat-Tyrant (*Ochthoeca rufipectoralis*) and House Wren (*Troglodytes aedon*) (Table 2). It is noteworthy that these bird species were mainly captured in high-altitude areas in Peru (i.e. Cuzco and Junin) where simuliids (vectors of *Leucocytozoon* species) can find optimal conditions, especially mountain streams, to complete their life-cycle (Greiner *et al.* 1975; Merino *et al.* 2008). As the Tamarugo Conebill migrates to high altitudes in southern Peru, it is likely that the three parasitised individuals were infected in these areas before

returning to their breeding regions in Pampa del Tamarugal where simuliids have not yet been reported.

Only House Sparrows were found to be infected with *Plasmodium* (prevalence 30%). The haplotype detected (GRW4) is thought to have been introduced to Hawaii by the House Sparrow, causing the subsequent decline and extinction of a number of endemic Hawaiian birds (van Riper 1991). This haplotype has been detected in several species of birds. It is a dominant haplotype on some oceanic islands, like Bermuda and French Polynesia (Beadell *et al.* 2006). In North America, it has thus far only been detected in House Sparrows, and has only rarely been detected in continental South America (Marzal *et al.* 2011). In fact, there are only eight reports, all from Argentina and Brazil and all isolated from House Sparrows (Marzal *et al.* 2011).

Therefore, it could be that the House Sparrow is spreading the parasite from north to south through the Americas. In addition, suitable vectors needed either to be present or introduced in parallel to facilitate parasite transmission in the regions invaded by the House Sparrow. In fact, the decline and extinction of endemic Hawaiian birds has been related to the introduction of *Culex pipiens quinquefasciatus* in 1827 (Zimmerman 1948a, 1948b). This vector has been reported in the village of Pica (González *et al.* 2012) where most of the House Sparrows were captured (9 of 10). It is possible that the life-cycle of *Plasmodium* haplotype GRW4 was predominantly urban but, even so, presents a risk of infection for the endemic Tamarugo Conebill because 37 of 89 (41.6%) captured were caught in the same area of Pica where most House Sparrows were captured. Although encounters between *Plasmodium* and Tamarugo Conebills is possible, the pathogenicity of the lineage GRW4 is not known. It is worth noting that previous studies have demonstrated the genetic diversity of the lineage GRW4 and its possible relationship to pathogenicity. In this sense, several TRAP (thrombospondin-related anonymous protein) and MSP1 (merozoite surface protein 1) alleles have been detected in GRW4 haplotypes isolated worldwide (Farias *et al.* 2012; Hellgren *et al.* 2015). These surface proteins are likely to interact with the immune system and parasite evasion mechanisms of the host. It is possible that parasites bearing different alleles behave as different populations whose transmission and pathogenic characteristics may be specific for a particular host species (Hellgren *et al.* 2015). Thus, it would be interesting to characterise the allelic diversity present in the haplotype GRW4 isolated in the present survey to establish its origin and potential pathogenicity.

The third haemosporidian parasite found in the Pampa del Tamarugal belongs to the subgenus *Haemoproteus*. It was only detected in the Eared Dove, with 100% prevalence ($n=4$). This parasite is transmitted by louse flies (family Hippoboscidae), which are obligate parasites and many are flightless. Most larval development takes place within the maternal body and pupation occurs almost immediately once in the soil or on the host. As parasites belonging to the subgenus *Haemoproteus* have not yet been isolated from passeriform birds (Santiago-Alarcon *et al.* 2014; Valkiūnas *et al.* 2013; Merino *et al.* 2012), those found in the Eared Dove should not represent a threat to the Tamarugo Conebill.

In conclusion, the overall prevalence of blood parasites in the present survey was low. However, haemoparasites transmitted by hippoboscids, such as *Haemoproteus multipigmentatus*, could be common in arid regions and overall prevalence may be higher with sufficient samples of scarcer species of birds. Further, although the main threat to Tamarugo Conebills relates to the management of the Tamarugo forests (BirdLife International 2012), the occurrence of House Sparrows infected with *Plasmodium* should be considered a risk. As noted above, a range of ecological factors might have contributed to the lack of past encounters between Tamarugo Conebills and the *Plasmodium* lineage isolated in the present study. As such, Tamarugo Conebills would not have acquired the immunological defence mechanisms to combat this parasite. To establish the actual risk of haemoparasites to the avian community, a more comprehensive survey of parasites and vectors over time, and the study of

any negative health effects of these parasites on birds, should be conducted.

Acknowledgements

During preparation of this work, J. Martínez and S. Merino were supported by funding from the Spanish Ministry of Science and Technology (project CGL2009–09439), the Spanish Ministry of Economy and Competitiveness (projects CGL2012–40026-C02–01 and 02) and the Fundación BBVA (project BIOCON06/109). Assistance with travel and accommodation to Pica was provided by the joint project between Consejo Superior de Investigaciones Científicas (CSIC, Spain) and Comisión Nacional de Investigación Científica y Tecnológica (CONICYT, Chile) (2009CL0025 / CONICYT-CSIC 2009–137). Research has been supported by Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT-Chile 1090794, FONDECYT 1140548) and the Institute of Ecology and Biodiversity (ICM-P05–002–Chile, and PFB-23–CONICYT–Chile), to R. A. Vásquez. We also acknowledge the support of A. Vilaxa, C. González and A. Taucare.

References

- Arenas, J. (2013). Observatorio del clima durante el año 2012 en la Estación Experimental de Canchones, pampa del Tamarugal, comuna de Pozo Almonte, provincia del Tamarugal, región de Tarapacá. Facultad de recursos naturales renovables, Universidad Arturo Prat, Chile.
- Beadell, J. S., Ishtiaq, F., Covas, R., Melo, M., Warren, B. H., Atkinson, C. T., Bensch, S., Graves, G. R., Jhala, Y. V., Peirce, M. A., Rahmani, A. R., Fonseca, D. M., and Fleischer, R. C. (2006). Global phylogeographic limits of Hawaii's avian malaria. *Proceedings of the Royal Society B: Biological Sciences* **273**, 2935–2944. doi:10.1098/rspb.2006.3671
- BirdLife International (2012) *Conirostrum tamarugense*. The IUCN Red List of Threatened Species 2012: e.T22722099A38171800. doi:10.2305/IUCN.UK.2012-1.RLTS.T22722099A38171800.en
- Clark, N. J., Clegg, S. M., and Lima, M. R. (2014). A review of global diversity in avian haemosporidians (*Plasmodium* and *Haemoproteus*: Haemosporida): new insights from molecular data. *International Journal for Parasitology* **44**, 329–338. doi:10.1016/j.ijpara.2014.01.004
- Combes, C. (2005). 'The Art of Being a Parasite.' (University Chicago Press: London.)
- Durrant, K. L., Beadell, J. S., Ishtiaq, F., Graves, G. R., Olson, S. L., Gering, E., Peirce, M. A., Milensky, C. M., Schmidt, B. K., Gebhard, C., and Fleischer, R. C. (2006). Avian hematozoa in South America: a comparison of temperate and tropical zones. *Ornithological Monographs* **60**, 98–111. doi:10.1642/0078-6594(2006)60[98:AHISAA]2.0.CO;2
- Estades, C. F. (1995). Aves de la reserva nacional Pampa del Tamarugal. *Boletín Chileno de Ornitología* **2**, 21–23.
- Estades, C. F. (1996). Natural history and conservation status of the Tamarugo Conebill in northern Chile. *Wilson Bulletin* **108**, 268–279.
- Farias, M. E. M., Atkinson, C. T., LaPointe, D. A., and Jarvi, S. I. (2012). Analysis of the *trap* gene provides evidence for the role of elevation and vector abundance in the genetic diversity of *Plasmodium relictum* in Hawaii. *Malaria Journal* **11**, 305. doi:10.1186/1475-2875-11-305
- González, C. R., Reyes, C., Rada, V., Jercic, M. I., Pavletic, C., and Parra, A. (2012). 'Manual de Culicidos (Diptera : Culicidae) de la Zona Norte y Centro de Chile, Incluyendo Isla de Pascua.' (Maval Impresores: Santiago, Chile.)
- González, A. D., Lotta, I. A., García, L. F., Moncada, L. I., and Matta, N. E. (2015). Avian haemosporidians from Neotropical highlands: evidence from morphological and molecular data. *Parasitology International* **64**, 48–59. doi:10.1016/j.parint.2015.01.007
- Greiner, E. C., Bennett, G. F., White, E. M., and Coombs, R. F. (1975). Distribution of the avian hematozoa of North America. *Canadian Journal of Zoology* **53**, 1762–1787. doi:10.1139/z75-211

- Hajek, E. R., and Di Castri, F. (1975). 'Bioclimatografía de Chile.' (Universidad Católica de Chile Press: Santiago, Chile.)
- Hellgren, O., Krizanauskienė, A., Hasselquist, D., and Bensch, S. (2011). Low haemosporidian diversity and one key-host species in a bird malaria community on a mid-Atlantic island (Sao Miguel, Azores). *Journal of Wildlife Diseases* **47**, 849–859. doi:10.7589/0090-3558-47.4.849
- Hellgren, O., Atkinson, C. T., Bensch, S., Albayrak, T., Dimitrov, D., Ewen, J. G., Kim, K. S., Lima, M. R., Martin, L., Palinauskas, V., Ricklefs, R., Sehgal, R. N. M., Valkiūnas, G., Tsuda, Y., and Marzal, A. (2015). Global phylogeography of the avian malaria pathogen *Plasmodium relictum* based on MSP1 allelic diversity. *Ecography* **38**, 842–850. doi:10.1111/ecog.01158
- LaPointe, D. A., Goff, M. L., and Atkinson, C. T. (2010). Thermal constraints to the sporogonic development and altitudinal distribution of avian malaria *Plasmodium relictum* in Hawai'i. *Journal of Parasitology* **96**, 318–324. doi:10.1645/GE-2290.1
- López-Calleja, M. V., and Estades, C. F. (1996). Natural history of the Tamarugo Conebill (*Conirostrum tamarugense*) during the breeding period: diet and habitat preferences. *Revista Chilena de Historia Natural (Valparaíso, Chile)* **69**, 351–356.
- Martínez, J., Martínez-de la Puente, J., Herrero, J., del Cerro, S., Lobato, E., Rivero-de Aguilar, J., Vásquez, R. A., and Merino, S. (2009). A restriction site to differentiate *Plasmodium* and *Haemoproteus* infections in birds: on the inefficiency of general primers for detection of mixed infections. *Parasitology* **136**, 713–722. doi:10.1017/S0031182009006118
- Martínez, J., Vásquez, R. A., Venegas, C., and Merino, S. (2015). Molecular characterisation of haemoparasites in forest birds from Robinson Crusoe Island: is the Austral Thrush a potential threat to endemic birds? *Bird Conservation International* **25**, 139–152. doi:10.1017/S0959270914000227
- Marzal, A., Ricklefs, R. E., Valkiūnas, G., Albayrak, T., Arriero, E., Bonneaud, C., Czirájk, G. A., Ewen, J., Hellgren, O., Horáková, D., Iezhova, T. A., Jensen, H., Krizanauskienė, A., Lima, R. M., de Lope, F., Magnussen, E., Martin, L. B., Möller, A. P., Palinauskas, V., Pap, P. L., Pérez-Tris, J., Sehgal, R. N. M., Soler, M., Szölloši, E., Westerdahl, H., Zetindjiev, P., and Bensch, S. (2011). Diversity, loss, and gain of malaria parasites in a globally invasive bird. *PLoS One* **6**(7), e21905. doi:10.1371/journal.pone.0021905
- Merino, S., and Potti, J. (1995). Mites and blowflies decrease growth and survival in nestling Pied Flycatchers. *Oikos* **73**, 95–103. doi:10.2307/3545730
- Merino, S., Potti, J., and Fargallo, J. A. (1997). Blood parasites of some passerine birds from central Spain. *Journal of Wildlife Diseases* **33**, 638–641. doi:10.7589/0090-3558-33.3.638
- Merino, S., Moreno, J., Sanz, J. J., and Arriero, E. (2000). Are avian blood parasites pathogenic in the wild? A medication experiment in Blue Tits. *Proceedings of the Royal Society B: Biological Sciences* **267**, 2507–2510. doi:10.1098/rspb.2000.1312
- Merino, S., Moreno, J., Vásquez, R. A., Martínez, J., Sánchez-Monsálvez, I., Estades, C. F., Ippi, S., Sabat, P., Rozzi, R., and Megehee, S. (2008). Haematozoa in forest birds from southern Chile: latitudinal gradients in prevalence and parasite lineage richness. *Austral Ecology* **33**, 329–340. doi:10.1111/j.1442-9993.2008.01820.x
- Merino, S., Hennicke, J., Martínez, J., Ludynia, K., Torres, R., Work, T. M., Stroud, S., Masello, J. F., and Quillfeldt, P. (2012). Infection by *Haemoproteus* parasites in four species of frigatebirds and the description of a new species of *Haemoproteus* (Haemosporida : Haemoproteidae). *Journal of Parasitology* **98**, 388–397. doi:10.1645/GE-2415.1
- Santiago-Alarcon, D., Rodríguez-Ferraro, A., Parker, P. G., and Ricklefs, R. E. (2014). Different meal, same flavor: cospeciation and host switching of haemosporidian parasites in some non-passerine birds. *Parasites & Vectors* **7**, 286. doi:10.1186/1756-3305-7-286
- Sehgal, R. N. M. (2015). Manifold habitat effects on the prevalence and diversity of avian blood parasites. *International Journal for Parasitology: Parasites and Wildlife* **4**, 421–430. doi:10.1016/j.ijppaw.2015.09.001
- Tompkins, D. M., and Gleeson, D. M. (2006). Relationship between avian malaria distribution and exotic invasive mosquito in New Zealand. *Journal of the Royal Society of New Zealand* **36**, 51–62. doi:10.1080/03014223.2006.9517799
- Valkiūnas, G. (2005). 'Avian Malaria Parasites and Other Haemosporidia.' (CRC Press: Boca Raton, FL.)
- Valkiūnas, G., Iezhova, T. A., Evans, E., Carlson, J. S., Martínez-Gómez, J. E., and Sehgal, R. N. M. (2013). Two new *Haemoproteus* species (Haemosporida : Haemoproteidae) from columbiform birds. *Journal of Parasitology* **99**, 513–521. doi:10.1645/12-98.1
- van Riper, C. III (1991). The impact of introduced vectors and avian malaria on insular passeriform bird populations in Hawaii. *Bulletin of the Society of Vector Ecology* **16**, 59–83.
- van Riper, C. III, van Riper, S. G., Goff, M. L., and Laird, M. (1986). The epizootiology and ecological significance of malaria in Hawaiian land birds. *Ecological Monographs* **56**, 327–344. doi:10.2307/1942550
- Watson, M. J. (2013). What drives population-level effects of parasites? Meta-analysis meets life-history. *International Journal for Parasitology: Parasites and Wildlife* **2**, 190–196. doi:10.1016/j.ijppaw.2013.05.001
- Zimmerman, E. C. (1948a). 'Insects of Hawaii. Vol. 1: Introduction.' (University of Hawaii Press: Honolulu.)
- Zimmerman, E. C. (1948b). 'Insects of Hawaii. Vol. 2: Apterigota to Thysanoptera.' (University of Hawaii Press: Honolulu.)